

# Asymptotic of sequences A161630, A212722, A212917 and A245265

(Václav Kotěšovec, published July 16 2014)

In the OEIS (On-Line Encyclopedia of Integer Sequences) published Paul D. Hanna in 2009 sequences [A161630](#), [A212722](#), [A212917](#) and I added in 2014 sequence [A245265](#), which can be generalized (for  $p \geq 1$ ) as the family of the sequences with an [exponential generating function](#)  $A(x)$ , satisfies functional equation

$$A(x) = \exp\left(\frac{x}{1 - x A(x)^p}\right)$$

or as the sum

$$a_n = \sum_{k=0}^n \frac{n!}{k!} * \binom{n-1}{n-k} * (1 + p * (n-k))^{k-1}$$

**Theorem** (V. Kotěšovec, July 15 2014):

Asymptotic

$$a_n \sim \frac{n^{n-1} p^{n-1+\frac{1}{p}} \left(1 + 2 * \text{LambertW}\left(\frac{\sqrt{p}}{2}\right)\right)^{n+\frac{1}{2}}}{e^n 2^{2*n+\frac{2}{p}} \text{LambertW}\left(\frac{\sqrt{p}}{2}\right)^{2*n+\frac{2}{p}-\frac{1}{2}} \sqrt{1 + \text{LambertW}\left(\frac{\sqrt{p}}{2}\right)}}$$

## Proof:

Following theorem by Edward A. Bender is (in case of implicit functions) very useful (for proof see [1], p.505 and also [4], p.469).

**Citation:** Edward A. Bender, "Asymptotic methods in enumeration" (1974), p.502, see [1]

**THEOREM 5.** Assume that the power series  $w(z) = \sum a_n z^n$  with nonnegative coefficients satisfies  $F(z, w) \equiv 0$ . Suppose there exist real numbers  $r > 0$  and  $s > a_0$  such that

- (i) for some  $\delta > 0$ ,  $F(z, w)$  is analytic whenever  $|z| < r + \delta$  and  $|w| < s + \delta$ ;
- (ii)  $F(r, s) = F_w(r, s) = 0$ ;
- (iii)  $F_z(r, s) \neq 0$ , and  $F_{ww}(r, s) \neq 0$ : and
- (iv) if  $|z| \leq r, |w| \leq s$ , and  $F(z, w) = F_w(z, w) = 0$ , then  $z = r$  and  $w = s$ .

Then

$$(7.1) \quad a_n \sim ((rF_z)/(2\pi F_{ww}))^{1/2} n^{-3/2} r^{-n},$$

where the partial derivatives  $F_z$  and  $F_{ww}$  are evaluated at  $z = r, w = s$ .

Bender's formula modified for [exponential generating function](#) is

$$\frac{a_n}{n!} \sim \frac{1}{n r^n} \sqrt{\frac{r F_z}{2\pi n F_{ww}}}$$

$$a_n \sim \frac{n^{n-1}}{e^n r^{n-1/2}} \sqrt{\frac{F_z}{F_{ww}}}$$

Now we have the implicit function

$$f(x, y) = e^{\frac{x}{1-xy^p}} - y$$

partial derivatives		
$F_z$	$\frac{\partial}{\partial x} f(x, y)$	$\frac{e^{\frac{x}{1-xy^p}}}{(xy^p - 1)^2}$
$F_w$	$\frac{\partial}{\partial y} f(x, y)$	$\frac{px^2y^{p-1}e^{\frac{x}{1-xy^p}}}{(xy^p - 1)^2} - 1$
$F_{ww}$	$\frac{\partial}{\partial y} \frac{\partial}{\partial y} f(x, y)$	$-\frac{px^2y^{p-2}e^{\frac{x}{1-xy^p}}(p(x^2y^p(y^p - 1) - 1) + (xy^p - 1)^2)}{(xy^p - 1)^4}$

r, s, are roots of the system of equations

$$e^{\frac{r}{1-rs^p}} - s = 0 \quad \frac{pr^2s^{p-1}e^{\frac{r}{1-rs^p}}}{(rs^p - 1)^2} - 1 = 0$$

From first equation we have

$$r = \frac{1}{s^p + \frac{1}{\log(s)}}$$

and second equation can be reduced

$$pr^2s^p = (rs^p - 1)^2$$

$$p s^p (\log(s))^2 = 1$$

$$s = e^{\frac{2W(\sqrt{p})}{p}} = \left( \frac{\sqrt{p}}{2W(\sqrt{p}/2)} \right)^{\frac{2}{p}}$$

where W is the [LambertW](#) function

$$r = \frac{4 \left( W(\sqrt{p}/2) \right)^2}{p \left( 1 + 2W(\sqrt{p}/2) \right)}$$


The asymptotic is then

$$a_n \sim \frac{n^{n-1}}{e^n r^{n-1/2}} \sqrt{\frac{F_z}{F_{ww}}} = e^{-n} n^{n-1} r^{-n} \sqrt{-\frac{s^{2-p}(rs^p - 1)^2}{pr(p(r^2s^p(s^p - 1) - 1) + (rs^p - 1)^2)}}$$

and after simplification

$$a_n \sim \frac{n^{n-1} p^{n-1+\frac{1}{p}} \left( 1 + 2 * W(\sqrt{p}/2) \right)^{\frac{1}{2}}}{e^n 2^{2*n+\frac{2}{p}} W(\sqrt{p}/2)^{2*n+\frac{2}{p}-\frac{1}{2}} \sqrt{1 + W(\sqrt{p}/2)}}$$

## Numerical verification (for p=4, sequence A245265)




---

## References:

- [1] Edward A. Bender, Asymptotic methods in enumeration, SIAM Review 16 (1974), no. 4, 485-515
- [2] Kotěšovec V., [Asymptotic of implicit functions if Fww = 0](#), extension of theorem by Bender, website 19.1.2014
- [3] [OEIS](#) - The On-Line Encyclopedia of Integer Sequences
- [4] P. Flajolet and R. Sedgewick, [Analytic Combinatorics](#), 2009, p. 469
- [5] Kotěšovec V., [Interesting asymptotic formulas for binomial sums](#), website 9.6.2013
- [6] Kotěšovec V., [Asymptotic of a sums of powers of binomial coefficients \\* x^k](#), website 20.9.2012
- [7] Kotěšovec V., [Asymptotic of sequences A244820, A244821 and A244822](#), website (and OEIS) 11.7.2014

This article was published on website <http://web.telecom.cz/vaclav.koteshovec/math.htm>, 16.7.2014  
and in the [OEIS](#), July 16 2014